



TITLE:

# On structure of CAT(0) groups (General and Geometric Topology today and their problems)

AUTHOR(S):

Hosaka, Tetsuya

---

CITATION:

Hosaka, Tetsuya. On structure of CAT(0) groups (General and Geometric Topology today and their problems). 数理解析研究所講究録 2013, 1833: 73-76

ISSUE DATE:

2013-05

URL:

<http://hdl.handle.net/2433/194866>

RIGHT:

## On structure of CAT(0) groups

静岡大学理学部

保坂 哲也 (Tetsuya Hosaka)

We introduce some results on finitely generated groups of isometries of CAT(0) spaces and CAT(0) groups in [15].

Definition and detail of CAT(0) spaces are found in [3] and [9].

Let  $X$  be a metric space and let  $\gamma$  be an isometry of  $X$ . Then the *translation length* of  $\gamma$  is defined as  $|\gamma| = \inf\{d(x, \gamma x) \mid x \in X\}$ , and the *minimal set* of  $\gamma$  is defined as  $\text{Min}(\gamma) = \{x \in X \mid d(x, \gamma x) = |\gamma|\}$ . An isometry  $\gamma$  is said to be *semi-simple* if  $\text{Min}(\gamma)$  is non-empty. Also an isometry  $\gamma$  is called

- (1) *elliptic* if  $\gamma$  has a fixed point,
- (2) *hyperbolic* if  $\gamma$  is semi-simple and not elliptic, and
- (3) *parabolic* if  $\gamma$  is not semi-simple.

(cf. [3, Chapter II.6]).

In [15], we obtained the following theorem by observing the proof of [3, Theorem II.6.12].

**Theorem 1.** *Let  $X$  be a CAT(0) space and let  $\Gamma$  be a finitely generated group acting by isometries on  $X$ . If the center of  $\Gamma$  contains a hyperbolic isometry  $\gamma_0$  of  $X$ , then there exist a normal subgroup  $\Gamma' \subset \Gamma$ , an element  $\delta_0 \in \Gamma$  and a number  $k_0 \in \mathbb{N}$  such that*

- (i)  $\Gamma = \Gamma' \rtimes \langle \delta_0 \rangle$ ,
- (ii)  $\Gamma' \rtimes \langle \delta_0^{k_0} \rangle = \Gamma' \times \langle \gamma_0 \rangle$  is a finite-index subgroup of  $\Gamma$  and
- (iii)  $\Gamma/\Gamma'$  is isomorphic to  $\mathbb{Z}$ .

A *geometric action* on a CAT(0) space is an action by isometries which is proper ([3, p.131]) and cocompact. A group  $\Gamma$  is called a *CAT(0) group*, if  $\Gamma$  acts geometrically on some CAT(0) space. We note that every CAT(0) space on

which some group acts geometrically is a proper space ([3, p.132]). Also we note that CAT(0) groups are finitely presented (cf. [3, Corollary I.8.11]).

For example, Bieberbach groups ([3, p.246], [4]), crystallographic groups ([4]), Coxeter groups and their torsion-free subgroups of finite-index ([6], [7], [19]) and fundamental groups of compact geodesic spaces of non-positive curvature ([3, p.159, p.237]) are CAT(0) groups. In particular, fundamental groups of Riemannian manifolds of non-positive sectional curvature are CAT(0) groups. Moreover, M. W. Davis [6] has constructed a closed aspherical manifold of dimension  $n \geq 5$  whose universal covering is not homeomorphic to  $\mathbb{R}^n$  ([6], [8]). The fundamental groups of these exotic manifolds are also CAT(0) groups.

On structure of CAT(0) groups, we obtained the following theorem from Theorem 1 in [15].

**Theorem 2.** *Let  $\Gamma$  be a CAT(0) group. Then there exist subgroups  $\Gamma = \Gamma_0 \supset \Gamma_1 \supset \cdots \supset \Gamma_n$ , elements  $\delta_{i+1}, \gamma_{i+1} \in \Gamma_i$  and  $k_{i+1} \in \mathbb{N}$  for  $i = 0, \dots, n-1$  such that*

- (1)  $\gamma_{i+1}$  is an element of the center of  $\Gamma_i$  with the order  $o(\gamma_{i+1}) = \infty$  for  $i = 0, \dots, n-1$ ,
- (2)  $\Gamma_i = \Gamma_{i+1} \rtimes \langle \delta_{i+1} \rangle$  for  $i = 0, \dots, n-1$ ,
- (3)  $\Gamma_{i+1} \rtimes \langle \delta_{i+1}^{k_{i+1}} \rangle = \Gamma_{i+1} \times \langle \gamma_{i+1} \rangle$  is a finite-index subgroup of  $\Gamma_i$ ,
- (4)  $\Gamma_i / \Gamma_{i+1}$  is isomorphic to  $\mathbb{Z}$  for  $i = 0, \dots, n-1$ ,
- (5)  $\Gamma = (\cdots ((\Gamma_n \rtimes \langle \delta_n \rangle) \rtimes \langle \delta_{n-1} \rangle) \rtimes \langle \delta_{n-2} \rangle) \cdots) \rtimes \langle \delta_1 \rangle$ ,
- (6)  $\Gamma_n$  has finite center, and
- (7)  $\Gamma_n \times A$  is a finite-index subgroup of  $\Gamma$  where  $A = \langle \gamma_1 \rangle \times \cdots \times \langle \gamma_n \rangle$  which is isomorphic to  $\mathbb{Z}^n$ .

Here, we introduce an easy example of a CAT(0) group.

**Example.** Let  $\Gamma = \langle a, b \mid ab^2 = b^2a \rangle$  and let  $X = \mathbb{R}^2$  the euclidean plane. We consider the action of the group  $\Gamma$  on  $X$  defined by

$$a \cdot (x, y) = (x, y + 1)$$

$$b \cdot (x, y) = (x + 1, -y)$$

for any  $(x, y) \in \mathbb{R}^2 = X$ . Then  $D = [0, 1] \times [-\frac{1}{2}, \frac{1}{2}] \subset \mathbb{R}^2$  is a fundamental domain,  $\Gamma D = X$  and  $\Gamma$  acts geometrically on  $X$ . Here we note that  $X/\Gamma$  is a

Klein bottle and the group  $\Gamma$  is a  $\text{CAT}(0)$  group which is the fundamental group of the Klein bottle.

Then  $\gamma_0 := b^2$  is a center of the  $\text{CAT}(0)$  group  $\Gamma$  and a hyperbolic isometry of  $X$ . Here we obtain that

- (i)  $\Gamma = \langle a \rangle \rtimes \langle b \rangle$ ,
- (ii)  $\langle a \rangle \rtimes \langle b^2 \rangle = \langle a \rangle \times \langle b^2 \rangle$  is a finite-index subgroup of  $\Gamma$  which is isomorphic to  $\mathbb{Z}^2$  and
- (iii)  $\Gamma / \langle a \rangle$  is isomorphic to  $\mathbb{Z}$ .

## REFERENCES

- [1] N. Bourbaki, *Groupes et Algèbres de Lie*, Chapters IV-VI, Masson, Paris, 1981.
- [2] P. Bowers and K. Ruane, *Boundaries of nonpositively curved groups of the form  $G \times \mathbb{Z}^n$* , Glasgow Math. J. **38** (1996), 177–189.
- [3] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [4] L. S. Charlap, *Bieberbach groups and flat manifolds*, Universitext. Springer-Verlag, New York, 1986.
- [5] C. B. Croke and B. Kleiner, *Spaces with nonpositive curvature and their ideal boundaries*, Topology **39** (2000), 549–556.
- [6] M. W. Davis, *Groups generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. **117** (1983), 293–324.
- [7] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.
- [8] M. W. Davis and T. Januszkiewicz, *Hyperbolization of polyhedra*, J. Diff. Geom. **34** (1991), 347–388.
- [9] E. Ghys and P. de la Harpe (ed), *Sur les groupes hyperboliques d'après Mikhael Gromov*, Progr. Math. vol. 83, Birkhäuser, Boston MA, 1990.
- [10] M. Gromov, *Hyperbolic groups*, Essays in group theory (S. M. Gersten, ed.), M.S.R.I. Publ. 8, 1987, pp. 75–264.
- [11] M. Gromov, *Asymptotic invariants for infinite groups*, Geometric Group Theory (G.A. Niblo and M.A. Roller, eds.), LMS Lecture Notes, vol. 182, Cambridge University Press, Cambridge, 1993, pp. 1–295.
- [12] P. de la Harpe, *Topics in geometric group theory*, Chicago Lectures in Math. Ser., The University of Chicago Press.
- [13] T. Hosaka, *Minimality of the boundary of a right-angled Coxeter system*, Proc. Amer. Math. Soc., **137** (2009), 899–910.
- [14] T. Hosaka, *On splitting theorems for  $\text{CAT}(0)$  spaces and compact geodesic spaces of non-positive curvature*, Math. Z., **272** (2012), 1037–1050.
- [15] T. Hosaka, *Remarks on structure of  $\text{CAT}(0)$  groups*, preprint.
- [16] J. E. Humphreys, *Reflection groups and Coxeter groups*, Cambridge University Press, 1990.
- [17] N. Monod, *Superrigidity for irreducible lattices and geometric splitting*, J. Amer. Math. Soc. **19** (2006), 781–814.
- [18] C. Mooney, *Examples of non-rigid  $\text{CAT}(0)$  groups from the category of knot groups*, Algebr. Geom. Topology **8** (2008), 1667–1690.

- [19] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, The Ohio State University, 1988.
- [20] J. M. Wilson, *A  $CAT(0)$  group with uncountably many distinct boundaries*, J. Group Theory 8 (2005), 229–238.